

Content available at: https://www.ipinnovative.com/open-access-journals

Indian Journal of Microbiology Research

Journal homepage: www.ijmronline.org



Original Research Article

Emergence of metronidazole and clindamycin resistance among anaerobic *Bacteroides fragilis* group - A study from South India

Sandra Jacob¹, Supriya Gachinmath¹*

¹Dept. of Microbiology, St. Johns Medical College. Bengaluru, Karnataka, India

Abstract

Background: *Bacteroides fragilis* is the most common pathogenic group among anaerobic pathogens, primarily causing infections such as deep visceral abscesses, brain abscesses, diabetic foot ulcers, etc. Recent studies have shown increasing resistance among these bacteria to common antibiotics like metronidazole and clindamycin, which are used for empirical treatment.

Aim and Objective: The study aimed to analyze and identify the common pathogenic species of the *Bacteroides fragilis* group isolated from clinical samples and to determine the drug resistance for clindamycin and metronidazole.

Materials and Methods: A total of 51 *Bacteroides fragilis* group isolates were obtained from samples received for anaerobic culture from January 2021 to December 2023. The samples were inoculated onto anaerobic blood agar and other essential media. Identification was done using MALDI-TOF MS. Antibiotic susceptibility was performed using E-strip method and interpreted according to CLSI 2023 breakpoints for clindamycin and metronidazole.

Results: *Bacteroides fragilis* was found to be the most prevalent among the *Bacteroides fragilis* group isolates, followed by *B. ovatus*. The cultures showed a high incidence of polymicrobial growth (80%, n=41), with *Prevotella* species (n=6) being the most frequent obligate anaerobe grown along with *Bacteroides fragilis* group, while *Escherichia coli* (n=15) was the most common facultative anaerobe. 11 out of 51 isolates showed resistance to metronidazole (21%), while 2 isolates were intermediate (3.9%) and the rest were sensitive. 23 out of 51 isolates (45%) were resistant to clindamycin, 1 was intermediate (1.9%), and 27 isolates were sensitive to clindamycin.

Conclusion: This study reiterates that *Bacteroides fragilis* subspp fragilis is the most common *Bacteroides fragilis* group and is known to cause various anaerobic infection. There is an emergence of drug resistance among them towards commonly used antibiotics like metronidazole and clindamycin.

Keywords: Bacteroides fragilis group, Obligate anaerobe, Antimicrobial resistance, Metronidazole, Clindamycin.

Received: 28-02-2025; Accepted: 07-09-2025; Available Online: 20-09-2025

This is an Open Access (OA) journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The *Bacteroides fragilis* group comprises gram-negative non-sporing obligate anaerobic bacilli. Similar to Enterobacterales, they are also part of the indigenous microbiota of the human colon. The Bacteroides genus has undergone many alterations since 1980s. Nowadays, the genus Bacteroides contains limited members of the *Bacteroides fragilis* group, including *Bacteroides fragilis*, which is the most common, *Bacteroides caccae*, *Bacteroides ovatus*, *Bacteroides eggerthii*, *Bacteroides thetaiotaomicron Bacteroides acidifaciens*, *Bacteroides coprocola*, *Bacteroides coprosuis*, *Bacteroides finegoldii*, etc.¹⁻³

They are opportunistic pathogens that primarily cause infection when they enter normal sterile sites in the body and are one of the most commonly isolated anaerobic bacteria in samples of human opportunistic infections. *A Bacteroides fragilis* group has been isolated in samples of various infections such as peritoneal abscess, brain abscess, diabetic foot ulcer, aspiration pneumonia, chronic suppurative otitis media, necrotizing fasciitis, pelvic abscess, etc. *5-8* They often cause infection in association with other anaerobic and aerobic bacteria. Isolation of anaerobic bacteria from clinical samples are difficult since they require special media (horse

*Corresponding author: Supriya Gachinmath Email: supriya.bg@stjohns.in blood agar, thioglycolate broth) along with specialized equipment for maintaining strict anaerobic conditions.9 Anaerobiosis can be achieved using various techniques like anaerobic chamber, anaerobic jars with Gaspak, anaerobic pouch systems, anoxomat etc. 9 Identification includes use of biochemical tests like Bacteroides bile esculin agar, use of special antibiotic disc like kanamycin(1000µg), vancomycin (5 μ g) and colistin (10 μ g) which are time consuming ⁹. Advances in the anaerobic culture techniques and usage of automated identification methods like Matrix-Assisted Laser Desorption/Ionization Time-of-Flight (MALDI-TOF) mass spectrometry have greatly improved the isolation and identification of these anaerobes, which was once cumbersome due to long incubation time, the need for special equipment, and high costs. This has helped us to understand the role of anaerobic bacteria like the Bacteroides fragilis group in various diseases, often associated with high degrees of morbidity and mortality.11 Routine antibiotic sensitivity tests are not done for anaerobic isolates in many clinical laboratories because the procedure requires special media and is often difficult and time-consuming.¹² Currently, metronidazole is used as an empirical antibiotic for anaerobic infections and prophylaxis.11

Studies conducted in the past two decades have shown an increasing trend of antibiotic resistance among anaerobic bacteria, with the highest incidence among the members of the *Bacteroides fragilis* group. ^{13,14} The resistance towards

clindamycin has been attributed to presence of *ermF* gene and metronidazole due to *nim* gene. 11,15 The need for routine testing of antibiotic susceptibility patterns has become crucial to mitigate incidences of antibiotic resistance. In this study, we have ascertained the antibiotic susceptibility of the *Bacteroides fragilis* group isolated from the various samples that were received at the microbiology laboratory of St. Johns medical college hospital in southern India.

2. Materials and Methods

The study was a descriptive cross-sectional study on Bacteroides fragilis group isolated from various samples received for anaerobic culture at the Department of Microbiology of St. Johns medical college hospital in Bangalore during the period from January 2021 to March 2024. Samples such as pus aspirates, body fluids, tissue from various sites suspected of anaerobic infection was cultured on anaerobic blood agar, Robertsons cooked meat broth, neomycin blood agar and brucella blood agar (Figure 1a). Additional tests such as growth on Bile esculin agar were also performed for confirmation of the genus Bacteroides (Figure 1b). The cultures that yielded Bacteroides fragilis group isolates were included in the study. Any swabs, like eye swabs, ear swabs, etc., were rejected, and fluid samples like urine samples and stool samples were not included in the study.




Figure 1: a): Bile Esculin agar with kanamycin showing the growth of Bacteroides fragilis (BF). **b**): Brucella blood agar showing the growth of translucent colonies of (BF). **c**): Anident disc sensitivity pattern of BF. **d**): E strip sensitivity of metronidazole showing MIC 0.75 which is sensitive

Anaerobiosis was achieved by either the anaerobic jar method with the commercially available gaspak (BD GasPakTM) or plastic GasPak pouch system (BD GasPakTM EZ Pouch Systems). Aerotolerance test was put up for each anaerobic isolate. Colonies that resembled the *Bacteroides fragilis* group were gram stained and colonies whose culture smears showed gram-negative bacilli resembling *Bacteroides fragilis* (**Figure 1**c) were identified using using Vitek MS (bioMérieux) which uses Matrix-assisted laser desorption/ionization-time of flight mass spectrometry (MALDI-TOF MS) technique.

Antibiotic susceptibility for metronidazole clindamycin was done by Epsilometer (E strip) method on Brucella Blood agar (Figure 1 d) using HiMedia Ezy MIC strip (EM019) for clindamycin and HiMedia Ezy MIC strip (EM 128) for metronidazole. The MIC was then compared to the CLSI 2023 breakpoints of the specific antibiotic for the Bacteroides fragilis group. 16 The isolate was identified as sensitive to Clindamycin if MIC was $\leq 2\mu g$, intermediate if MIC was 4µg, and resistant if MIC ≥8µg. ¹⁶ The isolate was identified as sensitive to Metronidazole if MIC was $\leq 4\mu g$, intermediate if MIC was $8\mu g$, and resistant if MIC $\geq 16\mu g$. ¹⁶ Data was analysed using SPSS v.28. All categorical data were summarized using frequency and percentages, and all continuous data were described using mean and standard deviation or Median based on the distribution.

3. Results

A total of 51 *Bacteroides fragilis* group isolates were isolated from various patient samples received at the Department of microbiology, St. Johns medical college. The majority of the patients were male (male to female ratio - 2.34:1). 54% of patients were above 45 years, 20 Patients were aged 16-44, 3 patients were less than 15 years of age, including 2 neonates, and the mean age was 46.9 years. *Bacteroides fragilis* (60%) was the common species isolated, followed by *Bacteroides ovatus* (19.6%) and *Bacteroides thetaiotaomicron* (11%) (**Figure 2**). *Bacteroides pyogenes, Bacteroides caccae*, and *Bacteroides stercoris* were the other species.




Figure 2: Species diversity of *Bacteroides fragilis* group grown in culture

The most common type of sample was tissue samples (47%), followed by pus aspirates (41%) and body fluids (8%)(**Figure 3**).




Figure 3: Showing the different samples from which *Bacteroides fragilis* group were isolated

In this study, 58% of the patients had co-morbidities, with Diabetes mellitus being the prevalent. Hypertension, thyroid disorders, and cardiac diseases were some of the other comorbidities associated with *Bacteroides fragilis* infection (**Table 1**).

Table 1: Various comorbidities associated with *Bacteroides fragilis* Group infections

Risk Factors (n=29)	Count		
Diabetes Mellitus	23		
Hypertension	16		
Thyroid Disorders	5		
Cardiac disease	7		
Other Comorbidities (Asthma,	bidities (Asthma, 5		
COPD, Paraplegia, etc)			

Most of the samples were obtained from the surgical ward / ICU, followed by the medical ward or ICU. The majority of the samples were from the gastrointestinal tract, followed by cutaneous tissue (**Table 2**). 80% of the samples were polymicrobial in growth (**Figure 4**), with *Prevotella* (n=6) being the most common obligate anaerobe grown along with the *Bacteroides fragilis* group.




Figure 4: Polymicrobial growth among the 51 samples

Other anaerobes include *Peptostreptococcus species* and *Finegoldia species*. The most common facultative anaerobe grown was *E.coli* (n=15) followed by *Streptococcus* species (n=8). Other organisms include *Enterococcus* (n=7) species and *Proteus* (n=3) species. Among Gram positive cocci isolated, *Enterococcus* (n=7) was the predominantly isolated. Most of the *Enterococcus* species (n=4) were Vancomycin

Resistant *Enterococcus*. 11 out of 51 *Bacteroides fragilis* group isolates showed resistance to metronidazole (21%), while 2 isolates were intermediate (3.9%) and the rest sensitive. 23 out of 51 isolates (45%) were resistant, 1 was intermediate (1.9%), and 26 isolates were sensitive to clindamycin. (**Figure 5**)




Figure 5: Antibiotic susceptibility of the Bacteroides fragilis group isolates

Table 2: The various clinical conditions from which Bacteroides fragilis group was isolated

S. No.	Systems involved	Conditions	No.
		Appendicular abscess	3
		Perianal abscess	2
		Abdominal wall abscess	4
	Gastrointestinal tract (22)	Peritoneal Abscess	4
1		Peri tonsillitis	2
1.		Pancreatitis	2
		Ischiorectal abscess	2
		Hirschsprung's disease	1
		Hepatolithiasis	1
		Esophago-pleural fistula	1
2	Subcutaneous and soft tissue (20)	Gangrene	6
		Fournier's gangrene	3
		Surgical site infection	2
		Bedsores	3
		Diabetic foot & ulcer	5
		Necrotizing fasciitis	1
3	Central nervous system (2)	VP Shunt Infection	1
		Cerebellar abscess	1
4	Genito urinary tract (1)	Fistula in the Genito urinary tract	1
5	Systemic infection (1)	Septic shock	1
6	Orofacial mandibular region (1)	Submandibular abscess	1
7	Other sites		4
	Total		51

4. Discussion

The *Bacteroides fragilis* group is one of the prevalent anaerobic pathogens infecting man and has been found to cause diseases ranging from abscess formation to sepsis. Often, in the case of suspected anaerobic infection, the patients are just given empirical antibiotic therapy with antibiotics like metronidazole and clindamycin. In the recent past, studies show an alarming incidence of drug resistance among the *Bacteroides fragilis* group, which has shown the importance of performing antibiotic susceptibility on anaerobic isolates.

In this study, the male-to-female patient ratio was found to be 2.34:1 among cultures showing growth of the *Bacteroides fragilis* group. Incidences of *Bacteroides fragilis* group infections were thus slightly more in samples of males than females. A similar male predominance was observed in the study conducted by Vishwanath S *et al.* in Manipal.¹¹ The most common age group affected was 61-75 years (27%) in our study.

Most patients were above the age of 45 years (n= 28, 54%), with 3 patients less than 15 years of age including 2 neonates. The study by Karlowsky *et al.* and others also states that the majority of the patients were also above 45 years of age. ^{11,17} Among the 51 patients whose samples showed significant growth of the *Bacteroides fragilis* group, 29 patients had one or more comorbidities. The most common comorbidity (**Table 1**) was found to be Type 2 Diabetes mellitus (79%) followed by Hypertension (55%).

The most common sample type (**Figure 3**) were tissue samples (47%), followed by pus aspirates (41%) and fluids (8%) unlike in the studies conducted by Lulu Jhan *et al.*, Vishwanath S *et al.*, and Jung Hyun Byun *et al.*, where the majority of samples were pus aspirates followed by tissue and body fluids. ^{11,15,18} Seventeen of the samples were aspirates from abscesses (33%), including one cerebellar abscess. This correlated with studies of Vishwanath and Nagy *et al.* ^{11,14}

Mostly, the samples were from the abdomen (**Table 2**), particularly the Gastrointestinal tract (43%). Samples from Gastrointestinal tract were the most common samples seen in the studies conducted by Lulu Jahan *et al.*, Vishwanath S *et al.*, E. Nagy *et al.*, and Fernández-Canigia *et al.*^{11,14,15,19} Other

infections include central nervous system infections (3.9%), gangrene (17.6%), and diabetic foot ulcers (9.8%).

A total of 41 (80%) out of the 51 samples showed polymicrobial growth of aerobic and anaerobic organisms, which is a characteristic of anaerobic infections (**Figure 4**). The studies conducted by Nakamura *et al.*, Elliott D *et al*, and Tzianabos *et al.* also conclude that most of the infections caused by the *Bacteroides fragilis* group are polymicrobial.²⁰⁻²² The common obligate anaerobes grown along with the *Bacteroides fragilis* group include *Prevotella species* (25%), followed by *Peptostreptococcus species* (13%) and Finegoldia species (13%) as seen in the studies conducted by Byun *et al.* and Irina *et al.* ^{18,23}

Bacteroides fragilis was found to be the most prevalent species (62.7%) isolated among the total 51 isolates of Bacteroides fragilis group followed by Bacteroides ovatus (17.3%) and Bacteroides thetaiotaomicron (11%). Other species isolated include Bacteroides vulgatus, Bacteroides pyogenes, and Bacteroides caccae (Figure 2). In the study conducted by James A. Karlowsky et al., Bacteroides fragilis followed by Bacteroides ovatus were the predominant species. Whereas in the study conducted by Viswanath et al., Nagy et al., and Lulu Jahan et al., Bacteroides fragilis followed by Bacteroides thetaiotaomicron were the commonly isolated species. 11,14,15

In this study, 11 out of 51 isolates showed resistance to metronidazole (21.5%), while 2 isolates were intermediate (3.9%) and 38 (74.5%) were sensitive to metronidazole. 23 out of 51 isolates (45%) were resistant, 1 intermediate (1.9%), and 27 (50%) isolates were sensitive to clindamycin. In a similar study conducted by Lulu Jahan *et al.* the resistance to metronidazole was found to be 12.5% and in the study by Viswananth S it was 7% of the isolates. Het of the isolates among *Bacteroides fragilis* group in a study conducted by Sethi S *et al.* in India. Where as in the studies conducted by Michael E. J. Buhl *et al.*, Viswananth S, Nagy E *et al.*, and James Karlowsky *et al.*, more than 20% *Bacteroides fragilis* group isolates were resistant to clindamycin. H1.14,17,25

The percentage of isolates showing resistance to metronidazole and clindamycin were significantly higher than previous studies (**Table 3**).

Table 3: Different studies on metronidazole and clindamycin resistance among Bacteroides fragilis group

Study	Year	Country/ region	% of Isolates resistant to metronidazole	% of Isolates resistant to Clindamycin
Michael E. J. Buhl et al. ²⁵	2024	Europe	1.8%	20.9%
Lulu Jahan <i>et al</i> . ¹⁵	2023	India	12.5%	-
Viswananth S et al.11	2019	India	7%	31%
Sethi S et al. ²⁴	2019	India	31%	-
Irina Shilnikova et al. ²³	2014	Russia	2.9%	22.4%
Fernández-Canigia et al. ¹⁹	2011	Argentina	0%	26%
Nagy E et al. ¹⁴	2010	Europe	1%	32%
James Karlowsky et al. ¹⁷	2011	Canada	<1%	48%

Escherichia coli (44%) and Streptococcus species (22%) were the common facultative anaerobes grown along with the Bacteroides fragilis group. Similar findings were seen in the study conducted by Elliot et al. and Viswanath et al. 11,22 About 45% of the facultative anaerobic GNB were resistant to meropenem, with 37% of Escherichia coli showing resistance, 50% Klebsiella spp and 50% Proteus spp showing resistance to meropenemin in our study. 95% of the GNB were resistant to ampicillin, cefotaxime, ceftazidime and cefuroxime, while almost 90% showed resistance to ciprofloxacin. Among Gram-positive cocci isolated, Enterococcus species (n=7) were the most common. Most of the Enterococcus species (n=4) were Vancomycin Resistant Enterococcus.

5. Conclusion

Bacteroides fragilis group pathogens are the most prevalent pathogenic anaerobe in clinical samples, but they are often overlooked in routine diagnostics due to the difficulties in isolation. This study shows the trend of an increase in resistance among the Bacteroides fragilis group towards metronidazole and clindamycin, which are the common antibiotics used to treat anaerobic infections. Hence, routine screening for the presence of anaerobic pathogens like the Bacteroides fragilis group is necessary, and empirical therapy should be followed by appropriate targeted antibiotic therapy for anaerobes after antibiotic susceptibility testing.

6. Limitations

No genotypic study was performed to access the presence of resistance genes like *nim* gene and *ermF* gene. The study was not multi centric limiting the findings to only a small population of the country. The resistance of *Bacteroides fragilis* group to other common antibiotics need to be assessed especially carbapenems.

7. Source of Funding

Nil.

8. Conflict of Interest

The authors declare that there is no conflict of interest.

9. Ethics Statement

The study was approved by the Internal ethics committee (IEC Ref no. TH-13/2023) of St. Johns Medical College, Bengaluru.

10. Consent and Confidentiality

Separate patient consent was not required as only the laboratory data on anaerobic cultures received as part of routine care was analysed in the study; the identity of the patient was not required or used in the data collection and analysis.

11. Authors' Contribution

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication. Jacob S was involved in the research, lab work, collection of data, interpretation, statistical analysis and maintenance of records. Gachinmath S was involved in the study topic, study design, data analysis and supervision.

12. Acknowledgements

Christobel Sobiya, Dept. of Microbiology. St Johns Medical College. Bengaluru.

References

- Wexler HM. Bacteroides: the good, the bad, and the nitty-gritty. Clin Microbiol Rev. 2007;20(4):593–621.
- Sakamoto M, Benno Y. Reclassification of Bacteroides distasonis, Bacteroides goldsteinii and Bacteroides merdae as Parabacteroides distasonis gen. nov., comb. nov., Parabacteroides goldsteinii comb. nov. and Parabacteroides merdae comb. nov. Int J Syst Evol Microbiol. 2006;56(Pt 7):1599–605.
- Hansen KCM, Schwensen SAF, Henriksen DP, Justesen US, Sydenham TV. Antimicrobial resistance in the *Bacteroides fragilis* group in faecal samples from patients receiving broad-spectrum antibiotics. *Anaerobe*. 2017;47:79–85.
- Nagy E. Anaerobic infections: update on treatment considerations. *Drugs*. 2010;70(7):841–58.
- Marik PE. Aspiration pneumonitis and aspiration pneumonia. N Engl J Med. 2001;344(9):665–71.
- Brook I. The role of anaerobic bacteria in chronic suppurative otitis media in children: Implications for medical therapy. *Anaerobe*. 2008;14(6):297–300.
- Green RJ, Dafoe DC, Rajfin TA. Necrotizing Fasciitis. Chest. 1996;110(1):219–29.
- Citron DM, Goldstein EJC, Merriam CV, Lipsky BA, Abramson MA. Bacteriology of moderate-to-severe diabetic foot infections and in vitro activity of antimicrobial agents. *J Clin Microbiol*. 2007;45(9):2819–28.
- Jousimies-Somer H, Summanen P, Citron D. Wadsworth-KTL anaerobic bacteriology manual. Belmont (CA): Start Now Press; 2002.
- Garner O, Mochon A, Branda J, Burnham C-A, Bythrow M, Ferraro M, et al. Multi-centre evaluation of mass spectrometric identification of anaerobic bacteria using the VITEK® MS system. Clin Microbiol Infect. 2014;20(4):335–9.
- 11. Shenoy P, Vishwanath S, Chawla K. Antimicrobial resistance profile and Nim gene detection among *Bacteroides fragilis* group isolates in a university hospital in South India. *J Glob Infect Dis*. 2019;11(2):59–62.
- Nagy E, Boyanova L, Justesen US. How to isolate, identify and determine antimicrobial susceptibility of anaerobic bacteria in routine laboratories. Clin Microbiol Infect. 2018;24(11):1139–48.
- 13. Yim J, Lee Y, Kim M, Seo YH, Kim WH, Yong D, et al. Antimicrobial susceptibility of clinical isolates of *Bacteroides fragilis* group organisms recovered from 2009 to 2012 in a Korean hospital. *Ann Lab Med*. 2015;35(1):94–8.
- Nagy E, Urbán E, Nord CE. Antimicrobial susceptibility of Bacteroides fragilis group isolates in Europe: 20 years of experience. Clin Microbiol Infect. 2011;17(3):371–9.
- Jahan L, Biswas R. Molecular study on metronidazole resistance in Bacteroides fragilis group isolates from a South Indian tertiary care center. Anaerobe. 2023;80:102692.
- CLSI. M100: Performance Standards for Antimicrobial Susceptability Testing; 2021.
- Karlowsky JA, Walkty AJ, Adam HJ, Baxter MR, Hoban DJ, Zhanel GG. Prevalence of antimicrobial resistance among clinical isolates of *Bacteroides fragilis* group in Canada in 2010-2011: CANWARD

- surveillance study. *Antimicrob Agents Chemother*. 2011;56(3):1247–52.
- Byun JH, Kim MS, Lee Y, Lee K, Chong Y. Antimicrobial susceptibility patterns of anaerobic bacterial clinical isolates from 2014 to 2016, including recently named or renamed species. *Ann Lab Med*. 2019;39(2):190–9.
- Fernández-Canigia L, Litterio M, Legaria MC, Prior C, Rossetti M, Vay C. First national survey of antibiotic susceptibility of the Bacteroides fragilis group: emerging resistance to carbapenems in Argentina. Antimicrob Agents Chemother. 2012;56(3):1309–14.
- Tzianabos AO, Onderdonk AB, Rosner B, Cisneros RL, Kasper DL. Structural features of polysaccharides that induce intra-abdominal abscesses. *Science*. 1993;262(5132):416–9.
- Nakamura I, Aoki K, Miura Y, Yamaguchi T, Matsumoto T. Fatal sepsis caused by multidrug-resistant *Bacteroides fragilis*, harboring a *cfiA* gene and an upstream insertion sequence element, in Japan. *Anaerobe*. 2017;44:36–9.
- Elliott D, Kufera JA, Myers RAM. The microbiology of necrotizing soft tissue infections. Am J Surg. 2000;179(5):361–6.

- Shilnikova II, Dmitrieva NV. Evaluation of antibiotic susceptibility of *Bacteroides*, *Prevotella* and *Fusobacterium* species isolated from patients of the N. N. Blokhin Cancer Research Center, Moscow, Russia. *Anaerobe*. 2015;31:15–8.
- Sethi S, Shukla R, Bala K, Gautam V, Angrup A, Ray P. Emerging metronidazole resistance in *Bacteroides* spp. and its association with the *nim* gene: a study from North India. *J Glob Antimicrob Resist*. 2019;16:210–4.
- Buhl MEJ, Sunnerhagen T, Join-Lambert O, Morris T, Jeverica S, Assous MV, et al. Antimicrobial resistance surveillance of Bacteroides fragilis isolated from blood cultures, Europe, 2022 (ReSuBacfrag). Int J Antimicrob Agents. 2024;64(3):107241.

Cite this article: Jacob S, Gachinmath S. Emergence of metronidazole and clindamycin resistance among anaerobic *Bacteroides fragilis* group - A study from South India. *Indian J Microbiol Res.* 2025;12(3):394–400.